

Infrared spectra of the Pd_nCO ($n=2-5$) molecules isolated in solid argon and neon between 100 and 4000 cm^{-1}

Benoît Tremblay, Sidi M. O. Souvi, Esmail Alikhani

- Sorbonne Université (Université Pierre et Marie Curie), Paris, France
- Laboratoire MONARIS: de la Molécule aux Nano-objets :

Réactivité, Interactions et Spectroscopies

THE Pd_nCO MOLECULES



- Important role played by the CO in transition metal complexes
- Model compounds to study the CO-palladium bond

THE Pd_nCO MOLECULES

- Important role played by the CO in transition metal complexes
- Model compounds to study the CO-palladium bond
- Pd_nCO : many theoretical works (~35)

THE Pd_nCO MOLECULES

- Important role played by the CO in transition metal complexes
- Model compounds to study the CO-palladium bond
- Pd_nCO : many theoretical works (~35), essentially for n= 2 and 3




THE Pd_nCO MOLECULES

- Important role played by the CO in transition metal complexes
 - Model compounds to study the CO-palladium bond
 - Pd_nCO : many theoretical works (~35), essentially for n= 2 and 3
 - Experimental results: CO absorbed on Pd particles or surface
- ⇒ CO absorbed on 1 metal atom: 2080-2100 cm⁻¹

THE Pd_nCO MOLECULES

- Important role played by the CO in transition metal complexes
- Model compounds to study the CO-palladium bond
- Pd_nCO : many theoretical works (~35), essentially for n= 2 and 3
- Experimental results: CO absorbed on Pd particles or surface
 - ⇒ CO absorbed on 1 metal atom: 2080-2100 cm⁻¹
 - ⇒ CO bridging 2 metal site: 1900-2000 cm⁻¹





THE Pd_nCO MOLECULES

- Important role played by the CO in transition metal complexes
- Model compounds to study the CO-palladium bond
- Pd_nCO : many theoretical works (~35), essentially for n= 2 and 3
- Experimental results: CO absorbed on Pd particles or surface

⇒ CO absorbed on 1 metal atom: 2080-2100 cm⁻¹

⇒ CO bridging 2 metal site: 1900-2000 cm⁻¹


⇒ CO on threefold-hollow site: 1800-1840 cm⁻¹




THE Pd_nCO MOLECULES

- Important role played by the CO in transition metal complexes
- Model compounds to study the CO-palladium bond
- Pd_nCO : many theoretical works (~35), essentially for n= 2 and 3
- Experimental results: CO absorbed on Pd particles or surface
- Matrix isolation is used to stabilize small metal molecules in quantities large enough to enable detection in the mid and far infrared
- Possibility to observe the low intensity bands


SAMPLE DEPOSITION



SAMPLE DEPOSITION




SAMPLE DEPOSITION




- Thermal evaporation: the metal atoms are in the ground state

SAMPLE DEPOSITION





- absorption spectra in the near-, mid- and far infrared were collected on the same sample, with a Bruker 120 FTIR spectrometer and InSb or MCT or bolometer detectors.

New bands in the CO stretching region




New bands in the CO stretching region




New bands in the CO stretching region

➡ Between 2040 and 2080 cm^{-1} : 1: m molecules




New bands in the CO stretching region


- In **argon** and neon : concentrations were varied over a large range
- Ne : easier atomic diffusion → higher formation of n: 1 with $n > 2$



New bands in the CO stretching region




New bands in the CO stretching region




New bands in the CO stretching region


$\text{PdCO}: \nu_{\text{C-O}} = 2044 \text{ cm}^{-1}$




New bands in the CO stretching region




New bands in the CO stretching region




New bands in the CO stretching region




New bands in the CO stretching region




New bands in the CO stretching region




New bands in the CO stretching region




all the bands were observed with $^{13}\text{C}^{16}\text{O}$ and $^{12}\text{C}^{18}\text{O}$ isotopes




New bands in the far infrared region




New bands in the far infrared region: argon



New bands in the far infrared region: argon




New bands in the far infrared region: argon




New bands in the far infrared region: argon

➡ all the bands were observed with $^{13}\text{C}^{16}\text{O}$ and $^{12}\text{C}^{18}\text{O}$ isotopes




New bands in the far infrared region: argon

Pd-C stretching mode




New bands in the far infrared region: argon


Pd-C stretching and **Pd-C-O bending modes**



Why two bands for the 2:1 molecule?




Photolysis: Irradiation at $\lambda = 405$ nm




No data on the electronic spectra of Pd_n

Only on atomic Pd

Photolysis: Irradiation at $\lambda = 405$ nm




Photolysis: Irradiation at $\lambda = 405$ nm



2:1 with end-on structure

2:1 with bridged structure

Photolysis: Irradiation at $\lambda = 405$ nm




2:1 with end-on structure

2015 cm^{-1}


2:1 with bridged structure

Photolysis: Irradiation at $\lambda = 405$ nm





From the calculations

➤ DFT with TPSS functional





From the calculations



2:1 with bridged structure is the most stable isomer in the singlet state


From the calculations





2:1 with bridged structure is the most stable isomer in the singlet state

Argon and neon

From the calculations




2:1 with end-on structure in the triplet state :
observed only in solid argon



2:1 with bridged structure is the most
stable isomer in the singlet state

From the calculations



2:1 with end-on structure in the triplet state :
observed only in solid argon

	²⁰ Ne	⁴⁰ Ar	⁸⁴ Kr	¹³¹ Xe	¹⁴ N ₂
Polarizability (10 ⁻²⁴ cm ³)	0.40	0.64	2.48	4.01	1.71


The higher polarizability of argon atoms stabilize this isomer

Comparison with the calculations

- DFT harmonic vibrational frequencies calculated with TPSS functional


Comparison with the calculations

Mode		Experiment			DFT-TPSS		
		¹² C ¹⁶ O	¹³ C ¹⁶ O	¹² C ¹⁸ O	¹² C ¹⁶ O	¹³ C ¹⁶ O	¹² C ¹⁸ O
2:1	γ_{PdCO}	387	-13	-4	369	-9	-2
br	δ_{OCPd}	588	-20	-2	575	-23	-3
	ν_{CO}	1857	-43	-41	1850	-41	-40
2:1	ν_{PdC}	484	-6	-14	454	-9	-14
eo	ν_{CO}	2015	-47	-44	2020	-46	-45






Comparison with the calculations

	Mode	Experiment			DFT-TPSS		
		¹² C ¹⁶ O	¹³ C ¹⁶ O	¹² C ¹⁸ O	¹² C ¹⁶ O	¹³ C ¹⁶ O	¹² C ¹⁸ O
2:1	γ_{PdCO}	387	-13	-4	369	-9	-2
br	δ_{OCPd}	588	-20	-2	575	-23	-3
	ν_{CO}	1857	-43	-41	1850	-41	-40
2:1	ν_{PdC}	484	-6	-14	454	-9	-14
eo	ν_{CO}	2015	-47	-44	2020	-46	-45
3:1	δ_{OCPd}	500	-17	-2	490	-17	-2
	ν_{CO}	1729	-39	-40	1730	-39	-40



Comparison with the calculations

Mode		Experiment			DFT-TPSS		
		¹² C ¹⁶ O	¹³ C ¹⁶ O	¹² C ¹⁸ O	¹² C ¹⁶ O	¹³ C ¹⁶ O	¹² C ¹⁸ O
2:1	γ_{PdCO}	387	-13	-4	369	-9	-2
br	δ_{OCPd}	588	-20	-2	575	-23	-3
	ν_{CO}	1857	-43	-41	1850	-41	-40
2:1	ν_{PdC}	484	-6	-14	454	-9	-14
eo	ν_{CO}	2015	-47	-44	2020	-46	-45
3:1	δ_{OCPd}	500	-17	-2	490	-17	-2
	ν_{CO}	1729	-39	-40	1730	-39	-40
4:1	δ_{OCPd}	515	-17	-2	521	-19	-1
	ν_{CO}	1694	-39	-40	1687	-38	-40




Comparison with the calculations


	Mode	Experiment			DFT-TPSS		
		$^{12}\text{C}^{16}\text{O}$	$^{13}\text{C}^{16}\text{O}$	$^{12}\text{C}^{18}\text{O}$	$^{12}\text{C}^{16}\text{O}$	$^{13}\text{C}^{16}\text{O}$	$^{12}\text{C}^{18}\text{O}$
2:1 br	γ_{PdCO}	387	-13	-4	369	-9	-2
	δ_{OCPd}	588	-20	-2	575	-23	-3
	ν_{CO}	1857	-43	-41	1850	-41	-40
2:1 eo	ν_{PdC}	484	-6	-14	454	-9	-14
	ν_{CO}	2015	-47	-44	2020	-46	-45
3:1	δ_{OCPd}	500	-17	-2	490	-17	-2
	ν_{CO}	1729	-39	-40	1730	-39	-40
4:1	δ_{OCPd}	515	-17	-2	521	-19	-1
	ν_{CO}	1694	-39	-40	1687	-38	-40

- The number of infrared active bands, their intensities, and their frequency positions provide insight into the structure and bonding of these complexes.


C-O frequency as a function of number of Pd atoms




C-O frequency as a function of number of Pd atoms




C-O frequency as a function of number of Pd atoms




C-O frequency as a function of number of Pd atoms



C-O frequency as a function of number of Pd atoms



C-O frequency as a function of number of Pd atoms



Conclusion

- Many new molecules have been experimentally detected

Conclusion

- Many new molecules have been experimentally detected
- Two Pd₂CO structures have been characterized in the mid- and far-infrared regions
- These two isomers can be interconnected by photochemistry

Conclusion

- Many new molecules have been experimentally detected
- Two Pd₂CO structures have been characterized in the mid- and far-infrared regions
- These two isomers can be interconnected by photochemistry
- The theoretical analysis of the data perfectly support the experimental results.

Thank you for your attention

